

# Avec®

(Cetirizine Dihydrochloride)

## Description:

### Each film coated tablet contains:

Cetirizine dihydrochloride (U.S.P.) equivalent to  
Cetirizine base 10 mg.  
Product Complies U.S.P. Specs.

### Each 5 ml contains:

Cetirizine dihydrochloride (U.S.P.) equivalent to  
Cetirizine base 5 mg.  
Product Complies Platinum Specs.

## Indications & Usage:

### Seasonal allergic rhinitis:

Symptoms treated effectively include sneezing, rhinorrhea, nasal pruritus, ocular pruritus, tearing, and redness of the eyes.

### Perennial allergic rhinitis:

Symptoms treated effectively include sneezing, rhinorrhea, post-nasal discharge, nasal pruritus, ocular pruritus, and tearing.

### Chronic urticaria

## Dosage & Administration:

### In adults and children over 12 years:

10 mg (1 tablet) O.D. or 5 mg (½ tablet) B.D.  
In patients with renal insufficiency dosage should be reduced to half a tablet daily.

### Children 6-12 years:

5 mg (1 teaspoon / 5 ml) or 10 mg (2 teaspoons / 10 ml) once daily depending on symptoms severity.

### Children 2-6 years:

2.5 mg (½ teaspoon / 2.5 ml) once or twice daily or  
5 mg (1 teaspoon / 5 ml) once daily or as directed by the physician.

## Precautions:

Though the drowsiness is very rare; due caution should therefore be exercised as it may affect the performance of skilled task like driving and operating machinery. Use in pregnancy and lactating women should be avoided.

## Drug Interaction:

As with other antihistamine it is advisable to avoid concomitant alcohol consumption.

## Side Effects:

Headache, dizziness, drowsiness, agitation, dry mouth and gastro-intestinal discomfort.

## Availability:

Avec film coated tablets are available in a blister pack of 10's.  
Avec syrup is available in apple flavour of 60 ml bottle.

## Warnings:

Store below 30°C, in a dry place, protect from light.  
Keep out of the reach of children.

# اویک

(سیٹی ریزین ڈائی ہائیڈروکلورائیڈ)

## خوراک:

بالغ اور ۱۲ سال سے بڑی عمر کے بچوں کے لئے:

۱۰ ملی گرام (ایک گولی) دن میں ایک مرتبہ یا

۵ ملی گرام (آدھی گولی) دن میں دو مرتبہ۔

۶ سال سے ۱۲ سال کے بچے:

ایک یا دو چائے کے چمچے دن میں ایک مرتبہ۔

۶ سال سے ۱۲ سال کے بچے:

آدھا چائے کا چمچے دن میں ایک یا دو مرتبہ یا

ایک چائے کا چمچے دن میں ایک مرتبہ یا

ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

○ دوا کو ۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر

روشنی سے بچا کر خشک جگہ پر رکھیں۔

○ تمام دوائیں بچوں کی پہنچ سے دور رکھیں۔

Manufactured by:

**Platinum**  
Pharmaceuticals (Pvt) Ltd.

A-20, North Western Industrial Zone,  
Bin Qasim, Karachi-75020, Pakistan.

QAR No. AW19-0397