

(Diclofenac Gel BP 1.16% w/w)

FASTAID GEL

Description:

Fastaid Gel contains:

Diclofenac diethylamine (BP)..... 1.16%w/w

equivalent to Diclofenac sodium.....1%w/w

Product Complies B.P Specs.

Fastaid Gel is a white, smooth gel without any lump or particles

Indications:

Fastaid gel is specially formulated for rubbing into the skin. It contains a substance called diclofenac which is one of the group of medicines called non-steroidal antiinflammatory analgesics, used to relieve pain and reduce swelling in a number of painful conditions affecting the joints and muscles.

Fastaid Gel relieves rheumatic and muscular pain, and reduces swelling and inflammation, e.g., in injuries involving the tendons, ligaments, muscles and joints. For the relief of sprains, strains or bruises, backache, tendinitis (e.g., tennis elbow), and for localised and mild arthritis.

Fastaid Gel may also be used to treat other conditions as determined by doctor.

ALLERGY AND CROSS-SENSITIVITY Contra-indicated in patients with a history of hypersensitivity to aspirin or any other NSAID-which includes those in whom attacks of asthma, angioedema, urticaria or rhinitis have been precipitated by aspirin or any other NSAID.

CAUTIONS Avoid contact with eyes, avoid contact with inflamed or broken skin, avoid contact with mucous membranes, not for use with occlusive dressings.topical application of large amounts can result in systemic effects, including hypersensitivity and asthma (renal disease has also been reported).

• **Pregnancy and Lactation:** Should consult the doctor before use.

SIDE EFFECTS:

Common or very common Conjunctivitis • muscle tone increased • oedema • rash (discontinue) • sensation abnormal • skin reactions

Uncommon Abdominal pain • alopecia • diarrhoea • eye pain • haemorrhage • lacrimation disorder • nausea • seborrhoea

Rare or very rare Asthma • hypersensitivity • photosensitivity reaction • rash pustular • renal failure

Frequency not known Hair colour changes

SIDE-EFFECTS, FURTHER INFORMATION Topical application of large amounts of diclofenac can result in systemic effects.

DIRECTIONS FOR ADMINISTRATION For topical preparations, apply with gentle massage only.

PATIENT AND CARER ADVICE For topical preparations, patients and their carers should be advised to wash hands immediately after use.

Storage conditions:

Store below 30°C in a dry place, protect from light and excessive heat avoid freezing.

Medicines should be kept out the reach of children,

Presentation:

Fastaid Gel is available in tube containing 20g gel, packed in carton with leaflet.

Manufactured by

Platinum

Pharmaceuticals (Pvt.) Ltd.

A-20 North Western Industrial Zone,

Bin Qasim, Karachi-75020, Pakistan.

URL: www.platinumpharma.net

QAR NO. AW20-0665-SLA

D2A

Platinum
Pharmaceuticals (Pvt.) Ltd.

Product : Fastaid Gel
Components: Leaflet
Country : Export (Srilanka)

Dimension: 170 mm x 145mm
Modified Date : 25-01-2021
Started Date: 22-05-2012
Concepts by : Sana
Design by : Rahil Rahim
QAR No. AW20-0665-SLA

Department	Marketing	PRO.	Regulatory	Quality Control	Quality Control Head	Procurement
Remarks						
Signature						
Date / Time						